

**To lower your risk of developing heart disease or stroke,are you...Eating fewer high fat or high cholesterol foods?
Kentucky BRFSS 2000**

Geographic Groups	Total Respond.#	Yes			No		
		N	%	C.I.(95%)	N	%	C.I.(95%)
TOTAL	6333	4154	65.2	(63.5-66.8)	2179	34.8	(33.2-36.5)
ADDs							
Barren River	436	281	63.9	(58.2-69.1)	155	36.1	(30.9-41.8)
Big Sandy	341	232	68.3	(62.6-73.6)	109	31.7	(26.4-37.4)
Bluegrass	497	332	65.9	(61.0-70.5)	165	34.1	(29.5-39.0)
Buffalo Trace	330	202	57.2	(50.9-63.3)	128	42.8	(36.7-49.1)
Cumberland Valley	394	248	60.2	(54.4-65.7)	146	39.8	(34.3-45.6)
FIVCO	459	308	61.8	(54.7-68.5)	151	38.2	(31.5-45.3)
Gateway	447	276	56.7	(51.0-62.2)	171	43.3	(37.8-49.0)
Green River	377	254	65.5	(59.8-70.8)	123	34.5	(29.2-40.2)
Kentucky River	422	293	67.4	(61.9-72.4)	129	32.6	(27.6-38.1)
KIPDA	468	327	70.3	(65.2-74.8)	141	29.7	(25.2-34.8)
Lake Cumberland	405	250	58.4	(52.5-64.0)	155	41.6	(36.0-47.5)
Lincoln Trail	432	275	61.4	(56.1-66.4)	157	38.6	(33.6-43.9)
Northern Kentucky	431	298	66.5	(61.3-71.4)	133	33.5	(28.6-38.7)
Pennyrile	428	280	62.4	(56.9-67.5)	148	37.6	(32.5-43.1)
Purchase	466	298	63.1	(57.9-68.0)	168	36.9	(32.0-42.1)

#Use caution in interpreting data for total respondents less than 50. N = Cell Size, NH = Non-Hispanic, % = Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level).

Percentages are weighted to population characteristics.

Unless stated differently above, Missing, 'Don't know', and Refused responses are excluded.